

Shoulder Arthroplasty

ORG: S-634 (ISC)

[Link to Codes](#)

MCG Health
Inpatient & Surgical
Care
27th Edition

- Care Planning - Inpatient Admission and Alternatives
 - Clinical Indications for Procedure
 - Alternatives to Procedure
 - Operative Status Criteria
 - Preoperative Care Planning
- Hospitalization
 - Optimal Recovery Course
 - Goal Length of Stay - **Ambulatory or 1 day postoperative**
 - Extended Stay
 - Hospital Care Planning
- Discharge
 - Discharge Planning
 - Discharge Destination
- Evidence Summary
 - Background
 - Criteria
 - Operative Status
 - Hospitalization
 - Length of Stay
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Care Planning - Inpatient Admission and Alternatives

Clinical Indications for Procedure

- Procedure is indicated for **1 or more** of the following(1)(2):
 - Advanced joint disease, as indicated by **ALL** of the following(3)(4):
 - Positive radiographic findings (eg, shoulder joint destruction, severe joint space narrowing, cystic changes)
 - Nonoperative therapy has been tried and failed (eg, anti-inflammatory medications, physical therapy, analgesics).
 - Replacement needed because of **1 or more** of the following:
 - Disabling pain
 - Functional disability
 - Osteonecrosis of the humeral head(5)(6)(7)
 - Reverse total shoulder arthroplasty, as indicated for **1 or more** of the following(8)(9):
 - Massive rotator cuff tear(10)
 - Failed rotator cuff repair(10)
 - Rotator cuff deficient arthropathy(10)
 - Failed shoulder arthroplasty or hemiarthroplasty(11)(12)
 - Proximal humerus fracture with rotator cuff deficiency or malunion(13)(14)(15)
 - Complex fracture of proximal humerus(14)(15)(16)(17)
 - Reconstruction after tumor resection(18)
 - Arthritis with posterior glenohumeral subluxation
 - Osteonecrosis of the humeral head(19)
 - Rheumatoid arthritis of the shoulder and **ALL** of the following(20)(21):
 - Positive radiographic findings (eg, shoulder joint destruction)
 - Nonoperative therapy has been tried and failed (eg, anti-inflammatory medications, disease-modifying antirheumatic drugs).
 - Replacement needed because of **1 or more** of the following:

- Disabling pain
- Functional disability
- ☐ Replacement (revision) of previous arthroplasty or hemiarthroplasty needed because of **1 or more** of the following(12):
 - Instability of glenoid or humeral components
 - Fracture or mechanical failure of implant
 - Glenoid erosion from humeral prosthetic component of hemiarthroplasty
 - Infection(22)(23)(24)(25)
 - Proximal migration of humeral head

Alternatives to Procedure

- Alternatives include(1)(4):
 - Nonoperative management, which may include(3)(10):
 - Anti-inflammatory medication
 - Analgesics
 - Flexibility and muscle strengthening exercises
 - Physical therapy
 - Intra-articular steroids
 - Reasonable restriction of activities
 - For proximal humerus fracture(13)(17)(26):
 - Open reduction. See Humerus Fracture, Closed or Open Reduction [ISC guideline](#).
 - Closed reduction with pin fixation or immobilization with orthosis
 - Nonoperative management(27)
 - Hemiarthroplasty. See Shoulder Hemiarthroplasty [ISC guideline](#).
 - Resection arthroplasty(28)
 - Radiation synovectomy
 - Arthroscopic debridement(29)
 - Serial debridement(22)(29)
 - Superior capsule reconstruction(30)(31)
 - Resurfacing of humeral head or glenoid(32)
 - Staged procedure with insertion of an antibiotic impregnated spacer(25)
 - Osteotomy for malunion
 - For osteonecrosis of humeral head(5)(6):
 - Humeral head core decompression
 - Humeral head bone grafting
 - Hemiarthroplasty. See Shoulder Hemiarthroplasty [ISC guideline](#).

Operative Status Criteria

- Ambulatory: Patients who are without active or unstable comorbidities (eg, CHF, COPD, CAD) that require prolonged postoperative care(33)(34)
- Inpatient: Patients with active or unstable comorbidities that require prolonged postoperative care(34)(35)[N](#)

Preoperative Care Planning

- Preoperative care planning needs may include(1):
 - Routine preoperative evaluation. See Preoperative Education, Assessment, and Planning Tool [SR](#).
 - Evaluation for metal hypersensitivity
 - Pain management(36)(37)
 - Preoperative treatment, procedures, and stabilization, including:
 - Ruling out sources of infection, including dental and lower urinary tract infections
 - Dental prophylaxis as indicated
 - Imaging
 - Preoperative discharge planning as appropriate. See Discharge Planning in this guideline.

Hospitalization

Optimal Recovery Course

Day	Level of Care	Clinical Status	Activity	Routes	Interventions	Medications
-----	---------------	-----------------	----------	--------	---------------	-------------

1	<ul style="list-style-type: none"> • OR to floor • Social Determinants of Health Assessment • Readmission Risk Assessment • Discharge planning • Possible discharge[A] 	<ul style="list-style-type: none"> • Clinical Indications met[B] 	<ul style="list-style-type: none"> • Ambulatory postoperatively • Possible pendulum exercises as tolerated 	<ul style="list-style-type: none"> • IV fluids • IV medications • Diet as tolerated postoperatively 	<ul style="list-style-type: none"> • Possible sling or shoulder immobilizer • Possible passive ROM postoperatively 	<ul style="list-style-type: none"> • Prophylactic antibiotics • Multimodal analgesia, possible PCA or continuous interscalene nerve block
2	<ul style="list-style-type: none"> • Social Determinants of Health Assessment • Readmission Risk Assessment • Floor to discharge • Complete discharge planning 	<ul style="list-style-type: none"> • Procedure completed • Hemodynamic stability • No evidence of neurologic or vascular compromise • No evidence of surgical site infection • Pain absent or managed • Discharge plans and education understood 	<ul style="list-style-type: none"> • Ambulatory or acceptable for next level of care • Pendulum exercises 	<ul style="list-style-type: none"> • Oral hydration[C] • Oral medications or regimen acceptable for next level of care • Oral diet or acceptable for next level of care 	<ul style="list-style-type: none"> • Physical therapy 	<ul style="list-style-type: none"> • PCA absent[D] • Possible analgesics

(1)(33)(34)NN

Recovery Milestones are indicated in **bold**.

Goal Length of Stay: Ambulatory or 1 day postoperative

Note: Goal Length of Stay assumes optimal recovery, decision making, and care. Patients may be discharged to a lower level of care (either later than or sooner than the goal) when it is appropriate for their clinical status and care needs.

Extended Stay




Minimal (a few hours to 1 day), Brief (1 to 3 days), Moderate (4 to 7 days), and Prolonged (more than 7 days).

- Extended stay beyond goal length of stay may be needed for(35)(38):
 - Complications of procedure(39)(40)(41)(42)
 - Complications include vascular or nerve injury, anterior dislocation, periprosthetic fracture, anesthetic complications, bleeding, and infection.
 - Expect brief stay extension.
 - Preoperative injury (eg, fracture, rotator cuff tear, multiple trauma)
 - Expect brief stay extension.
 - Care for comorbidities (eg, anemia, renal insufficiency)(43)(44)(45)(46)
 - Expect brief stay extension.

See Common Complications and Conditions [↗](#) ISC for further information.








Hospital Care Planning

- Hospital evaluation and care needs may include(1)(38):[N](#)
 - Treatment and procedure scheduling and completion, including:
 - Venous thromboembolism prophylaxis(51)
 - IV antibiotics

- Tranexamic acid(52)
- Transfusion
- Pain management(37)
- Consultation, assessment, and other services scheduling and completion, including:
 - Physical therapy(53)
- Identification of patient at high risk for readmission to prioritize transition and post-acute care
 - ☐ Risk of readmission is increased by presence of **1 or more** of the following(54)(55)(56)(57)(58)(59)(60)(61)(62)(63):
 - Hospitalization (nonelective) in past 6 months(64)(65)(66)
 - 2 or more emergency department visits in past 6 months
 - No source of outpatient care other than emergency department (eg, no primary care provider)(66)(67)
 - Severe care transition barriers (eg, no caregiver, homeless)(64)(65)(68)
 - Severe or end-stage renal disease (on dialysis or GFR less than 30 mL/min/1.73m² (0.5 mL/sec/1.73m²))(64)(69)
 -  eGFR - Adult Calculator  eGFR - Pediatric Calculator
 - Anemia(50)
 - Revision shoulder arthroplasty(47)
 - Inflammatory arthritis etiology(48)
 - Diabetes mellitus necessitating treatment with insulin(49)
 - Arthroplasty performed due to traumatic proximal humerus fracture(16)
- Monitoring patient's status for deterioration and comorbid conditions (see Inpatient Monitoring and Assessment Tool  SR); key items include:
 - Neurovascular status
 - Wound management
 - Pain management

Discharge

Discharge Planning

- Discharge planning includes[E]:
 - Assessment of needs and planning for care, including(71):
 - Develop treatment plan (involving multiple providers as needed).
 - Evaluate and address preadmission functioning as needed.
 - Evaluate and address psychosocial status issues as indicated. See Psychosocial Assessment  SR for further information.
 - Evaluate and address social determinants of health (eg, housing, food). See Social Determinants of Health Screening Tool  SR for further information.(70)
 - Evaluate and address patient or caregiver preferences as indicated.
 - Identify skilled services needed at next level of care, with specific attention to(72):
 - Neurovascular status assessment(73)
 - Pain management
 - Wound or dressing management
 - Early identification of anticipated discharge destination; options include(74)(75):
 - Home, considerations include:
 - Access to follow-up care
 - Home safety assessment. See Home Safety Assessment  SR for further information.
 - Self-management ability if appropriate. See Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment  SR for further information.
 - Caregiver need, ability, and availability
 - Post-acute skilled care or custodial care as indicated. See Discharge Planning Tool  SR for further information.
 - Transitions of care plan complete, including(75):
 - Patient and caregiver education complete. See Shoulder Arthroplasty: Patient Education for Clinicians  SR for further information.
 - See Teach Back Tool  SR for further information.
- ☐ Medication reconciliation completion includes(76)(77):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.
 - Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.

- Confirm that patient or caregiver can acquire medication.
- Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(78)
- Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
- See Medication Reconciliation Tool [↗](#) SR for further information.
- Plan communicated to patient, caregiver, and all members of care team, including(79)(80):
 - Inpatient care and service providers
 - Primary care provider
 - All post-discharge care and service providers
- Appointments planned or scheduled, which may include:
 - Primary care provider
 - Orthopedic surgeon(81)
 - Rehabilitation therapy services(1)(53)
 - Specialists for management of comorbidities as needed(81)
 - Other
- Outpatient testing and procedure plans made, which may include:
 - Other
- Referrals made for assistance or support, which may include:
 - Financial, for follow-up care, medication, and transportation
 - Tobacco use treatment(82)
 - Other
- Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), which may include:
 - Immobilizers (eg, braces, splints)(83)
 - Wound care equipment and supplies(84)
 - Other

Discharge Destination

- Post-hospital levels of admission may include:
 - Home.
 - Home healthcare. See Home Care Indications for Admission Section [↗](#) HC in Shoulder Arthroplasty guideline in Home Care.
 - Recovery facility care. See Recovery Facility Care Indications for Admission Section [↗](#) RFC in Shoulder Arthroplasty guideline in Recovery Facility Care.

Evidence Summary

Background

A shoulder arthroplasty involves replacement of the ball and socket and may include utilizing components which are opposite of the normal anatomy of the ball and socket (known as a reverse total shoulder arthroplasty). This guideline should be used for total shoulder arthroplasty, reverse total shoulder arthroplasty, and revision shoulder arthroplasty.(1) **(EG 2)**

Criteria

The evidence for the clinical indications found in this guideline includes 22 published peer reviewed articles, 2 specialty society or other evidence-based guidelines, and 1 book section.

Operative Status

Patients who are without active or unstable comorbidities (eg, CHF, COPD, CAD) that require prolonged postoperative care can be treated on an outpatient basis.(33)(34) **(EG 2)** Patients with active or unstable comorbidities that require prolonged postoperative care are treated on an inpatient basis.(34)(35) **(EG 3)**

Hospitalization

Readmission risk and reduction: Multivariate analysis of a database including 3501 patients (70% older than 65 years) who underwent shoulder arthroplasty found that having a revision arthroplasty procedure was independently associated with an increased risk of 30-day readmission.(47) **(EG 2)** Database analysis of 5801 patients who underwent total shoulder arthroplasty found, after multivariate adjustment, that inflammatory arthritis etiology (as opposed to osteoarthritis) was an independent risk factor for readmission at 30 days.(48) **(EG 2)** Database analysis of total shoulder arthroplasties (7246 patients) found, after multivariate adjustment, that the

presence of diabetes treated with insulin was independently associated with an increased risk for 30-day readmission.(49) **(EG 2)** Database analysis of total shoulder arthroplasties performed for degenerative osteoarthritis or traumatic proximal humeral fractures (8950 patients) found, after multivariate analysis, that total shoulder arthroplasty performance due to traumatic proximal humerus fractures was independently associated with increased risk of 30-day readmission.(16) **(EG 2)** Multivariate database analysis of 15,185 patients who underwent primary total shoulder arthroplasty found that preoperative anemia was independently associated with an increased risk of readmission within 30 days.(50) **(EG 2)**

Length of Stay

Analysis of a cohort of 370 consecutive total shoulder arthroplasty patients (mean age 71 years) found that 60% were discharged the day of surgery.(33) **(EG 2)**

Patients discharged the day after surgery can be considered inpatients or outpatients, depending on regulation or contractual arrangements. Whether considered inpatients or outpatients, analysis of data for a commercially insured population shows 90% of patients undergoing a principal procedure of shoulder arthroplasty were discharged the day of or the day after surgery.(34) **(EG 3)** A similar analysis for Medicare-insured patients shows 73% of patients were discharged the day of or the day after surgery.(34) **(EG 3)**

Rationale

Surgical MCG care guidelines help the clinician to identify, for a given procedure, which patient-specific factors and clinical conditions are appropriate for that procedure. The evidence-based clinical indication criteria assist the clinician in the decision to appropriately perform a procedure, evaluating whether the potential benefits of a procedure outweigh the potential risks. For Medicare enrollees, surgical MCG care guidelines also identify which procedures CMS has designated as inpatient only.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 412.3(85); 42 CFR 419.22(n)(86); 42 CFR 422.101(87)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(88); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 6 - Hospital Services Covered Under Part B(89); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(90); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment(91)

Medicare Coverage Determinations: Medicare Coverage Database(92)

References

1. Throckmorton TW. Shoulder and elbow arthroplasty. In: Azar FM, Beaty JH, editors. Campbell's Operative Orthopaedics. 14th ed. Philadelphia, PA: Elsevier; 2021:600-655.e8. [Context Link 1, 2, 3, 4, 5, 6, 7]
2. Le Breton S, et al. A validated algorithm using current literature to judge the appropriateness of anatomic total shoulder arthroplasty utilizing the RAND/UCLA appropriateness method. *Journal of Shoulder and Elbow Surgery* 2022;31(7):e332-e345. DOI: 10.1016/j.jse.2021.12.025. [Context Link 1] View abstract...
3. American Academy of Orthopaedic Surgeons. Management of Glenohumeral Joint Osteoarthritis. Evidence-based Clinical Practice Guideline [Internet] American Academy of Orthopaedic Surgeons. 2020 Mar Accessed at: <https://www.aaos.org/>. [created 2009; accessed 2022 Oct 10] [Context Link 1, 2]
4. Getz CL, Ricchetti ET, Verborgt O, Brolin TJ. Normal and pathoanatomy of the arthritic shoulder: considerations for shoulder arthroplasty. *Journal of the American Academy of Orthopedic Surgeons* 2019;27(24):e1068-e1076. DOI: 10.5435/JAAOS-D-18-00414. [Context Link 1, 2] View abstract...
5. Alkhateeb JM, Arafah MA, Tashkandi M, Al Qahtani SM. Surgical treatment of humeral head avascular necrosis in patients with sickle cell disease: a systematic review. *JSES International* 2021;5(3):391-397. DOI: 10.1016/j.jseint.2021.01.011. [Context Link 1, 2] View abstract...
6. Ristow JJ, et al. Outcomes of shoulder replacement in humeral head avascular necrosis. *Journal of Shoulder and Elbow Surgery* 2019;28(1):9-14. DOI: 10.1016/j.jse.2018.06.031. [Context Link 1, 2] View abstract...
7. Ayyash AM, Bui TM, O'Brien MJ, Mulcahey MK. Management of posttraumatic avascular necrosis of the proximal humerus. *Orthopedics* 2021;44(6):367-375. DOI: 10.3928/01477447-20211001-02. [Context Link 1] View abstract...
8. Almasri M, Kohrs B, Fleckenstein CM, Nolan J, Wendt A, Hasan SS. Reverse shoulder arthroplasty in patients 85 years and older is effective, safe, and durable. *Journal of Shoulder and Elbow Surgery* 2022;31(11):2287-2297. DOI: 10.1016/j.jse.2022.03.024. [Context Link 1] View abstract...
9. Marigi EM, et al. Reverse shoulder arthroplasty after prior rotator cuff repair: a matched cohort analysis. *Journal of the American Academy of Orthopedic Surgeons* 2022;30(3):e395- e404. DOI: 10.5435/JAAOS-D-21-00543. [Context Link 1] View abstract...
10. Cvetanovich GL, Waterman BR, Verma NN, Romeo AA. Management of the irreparable rotator cuff tear. *Journal of the American Academy of Orthopedic Surgeons* 2019;27(24):909-917. DOI: 10.5435/JAAOS-D-18-00199. [Context Link 1, 2, 3, 4] View abstract...

11. Harrison AK, Knudsen ML, Braman JP. Hemiarthroplasty and total shoulder arthroplasty conversion to reverse total shoulder arthroplasty. *Current Reviews in Musculoskeletal Medicine* 2020;13(4):501-508. DOI: 10.1007/s12178-020-09649-5. [Context Link 1] View abstract...
12. Weatherby PJ, Efejuku TA, Somerson JS. Complications after anatomic shoulder arthroplasty: revisiting leading causes of failure. *Orthopedic Clinics of North America* 2021;52(3):269-277. DOI: 10.1016/j.ocl.2021.03.002. [Context Link 1, 2] View abstract...
13. Kelly BJ, Myeroff CM. Reverse shoulder arthroplasty for proximal humerus fracture. *Current Reviews in Musculoskeletal Medicine* 2020;13(2):186-199. DOI: 10.1007/s12178-020-09597-0. [Context Link 1, 2] View abstract...
14. Boin MA, Virk MS. CORR® Synthesis: what is the role of reverse shoulder arthroplasty for the treatment of proximal humerus fractures in patients older than 65 years? *Clinical Orthopaedics and Related Research* 2021;479(11):2421-2429. DOI: 10.1097/CORR.0000000000001910. [Context Link 1, 2] View abstract...
15. Davey MS, et al. Management options for proximal humerus fractures - A systematic review & network meta-analysis of randomized control trials. *Injury* 2022;53(2):244-249. DOI: 10.1016/j.injury.2021.12.022. [Context Link 1, 2] View abstract...
16. Malik AT, Bishop JY, Neviasser AS, Beals CT, Jain N, Khan SN. Shoulder arthroplasty for a fracture is not the same as shoulder arthroplasty for osteoarthritis: implications for a bundled payment model. *Journal of the American Academy of Orthopedic Surgeons* 2019;27(24):927-932. DOI: 10.5435/JAAOS-D-18-00268. [Context Link 1, 2, 3] View abstract...
17. Mease SJ, Kraeutler MJ, Gonzales-Luna DC, Gregory JM, Gardner MJ, Choo AM. Current controversies in the treatment of geriatric proximal humeral fractures. *Journal of Bone and Joint Surgery. American Volume* 2021;103(9):829-836. DOI: 10.2106/JBJS.20.00665. [Context Link 1, 2] View abstract...
18. Ferlauto HR, et al. Reverse total shoulder arthroplasty for oncologic reconstruction of the proximal humerus: a systematic review. *Journal of Shoulder and Elbow Surgery* 2021;30(11):Online. DOI: 10.1016/j.jse.2021.06.004. [Context Link 1] View abstract...
19. McLaughlin R, et al. Reverse shoulder arthroplasty yields similar results to anatomic total shoulder arthroplasty for the treatment of humeral head avascular necrosis. *Journal of Shoulder and Elbow Surgery* 2022;31(6S):S94-S102. DOI: 10.1016/j.jse.2021.11.011. [Context Link 1] View abstract...
20. Leroux TS, Basques BA, Saltzman BM, Nicholson GP, Romeo AA, Verma NN. Shoulder arthroplasty in patients with rheumatoid arthritis: a population-based study examining utilization, adverse events, length of stay, and cost. *American Journal of Orthopedics (Belle Mead, N.J.)* 2018;47(6):Online. DOI: 10.12788/ajo.2018.0046. [Context Link 1] View abstract...
21. Haleem A, Shanmugaraj A, Horner NS, Leroux T, Khan M, Alolabi B. Anatomic total shoulder arthroplasty in rheumatoid arthritis: A systematic review. *Shoulder and Elbow* 2022;14(2):142-149. DOI: 10.1177/1758573220954157. [Context Link 1] View abstract...
22. Osmon DR, et al. Diagnosis and management of prosthetic joint infection: clinical practice guidelines by the Infectious Diseases Society of America. *Clinical Infectious Diseases* 2013;56(1):e1-e25. DOI: 10.1093/cid/cis803. (Reaffirmed 2022 Jun) [Context Link 1, 2] View abstract...
23. Garrigues GE, Zmistowski B, Cooper AM, Green A, ICM Shoulder Group. Proceedings from the 2018 International Consensus Meeting on Orthopedic Infections: management of periprosthetic shoulder infection. *Journal of Shoulder and Elbow Surgery* 2019;28(6S):S67-S99. DOI: 10.1016/j.jse.2019.04.015. [Context Link 1] View abstract...
24. Gross CE, Della Valle CJ, Rex JC, Traven SA, Durante EC. Fungal periprosthetic joint infection: a review of demographics and management. *Journal of Arthroplasty* 2021;36(5):1758-1764. DOI: 10.1016/j.arth.2020.11.005. [Context Link 1] View abstract...
25. Ruditsky A, McBeth Z, Curry EJ, Cusano A, Galvin JW, Li X. One versus 2-stage revision for shoulder arthroplasty infections: a systematic review and analysis of treatment selection bias. *Journal of Bone and Joint Surgery Reviews* 2021;9(9):Online. DOI: 10.2106/JBJS.RVW.20.00219. [Context Link 1, 2] View abstract...
26. London DA, et al. Impact of increasing comorbidity burden on resource utilization in patients with proximal humerus fractures. *Journal of the American Academy of Orthopedic Surgeons* 2020;28(21):e954-e961. DOI: 10.5435/JAAOS-D-19-00491. [Context Link 1] View abstract...
27. Handoll HH, Elliott J, Thillemann TM, Aluko P, Brorson S. Interventions for treating proximal humeral fractures in adults. *Cochrane Database of Systematic Reviews* 2022, Issue 6. Art. No.: CD000434. DOI: 10.1002/14651858.CD000434.pub5. [Context Link 1] View abstract...
28. George DA, Volpin A, Scarponi S, Haddad FS, Romano CL. Does exchange arthroplasty of an infected shoulder prosthesis provide better eradication rate and better functional outcome, compared to a permanent spacer or resection arthroplasty? a systematic review. *BMC Musculoskeletal Disorders* 2016;17:52. DOI: 10.1186/s12891-016-0901-6. [Context Link 1] View abstract...
29. Acosta-Olivo C, Vilchez-Cavazos F, Blazquez-Saldana J, Villarreal-Villarreal G, Pena-Martinez V, Simental-Mendia M. Comparison of open arthrotomy versus arthroscopic surgery for the treatment of septic arthritis in adults: a systematic review and meta-analysis. *International Orthopaedics* 2021;45(8):1947-1959. DOI: 10.1007/s00264-021-05056-8. [Context Link 1, 2] View abstract...
30. Eppler MB, et al. Superior capsular reconstruction of the shoulder. *Arthroscopy* 2021;37(6):1708-1710. DOI: 10.1016/j.arthro.2021.04.001. [Context Link 1] View abstract...
31. Werthel JD, et al. Superior capsular reconstruction - A systematic review and meta-analysis. *Orthopaedics & Traumatology, Surgery & Research : OTSR* 2021;107(8S):Online. DOI: 10.1016/j.otsr.2021.103072. [Context Link 1] View abstract...
32. Chillemi C, Paglialonga C, De Giorgi G, Proietti R, Carli S, Damo M. Outcome and revision rate of uncemented humeral head resurfacing: Mid-term follow-up study. *World Journal of Orthopedics* 2021;12(6):403-411. DOI: 10.5312/wjo.v12.i6.403. [Context Link 1] View abstract...
33. Cannon D, Lewis S, Garcia J, Watkins A, Rodriguez H, Levy JC. A Comparison of patient same day discharge (SDD) selection following shoulder arthroplasty before and after the COVID-19 pandemic. *Seminars in Arthroplasty* 2022;32(3):559-563. DOI: 10.1053/j.sart.2022.02.011. [Context Link 1, 2, 3, 4] View abstract...
34. Proprietary health insurance data sources (2020-2021); and Medicare 5% Standard Analytical File (2019-2020). [Context Link 1, 2, 3, 4, 5, 6, 7]
35. Premier PINC AI™ Healthcare Database (PHD), 01/01/2020-12/31/2021. Premier, Inc. [Context Link 1, 2, 3]
36. Codding JL, Getz CL. Pain management strategies in shoulder arthroplasty. *Orthopedic Clinics of North America* 2018;49(1):81-91. DOI: 10.1016/j.ocl.2017.08.010. [Context Link 1] View abstract...

37. Xiao M, Cohen SA, Cheung EV, Freehill MT, Abrams GD. Pain management in shoulder arthroplasty: a systematic review and network meta-analysis of randomized controlled trials. *Journal of Shoulder and Elbow Surgery* 2021;30(11):2638-2647. DOI: 10.1016/j.jse.2021.06.008. [Context Link 1, 2] View abstract...
38. Anakwenze OA, O'Donnell EA, Jobin CM, Levine WN, Ahmad CS. Medical complications and outcomes after total shoulder arthroplasty: a nationwide analysis. *American Journal of Orthopedics (Belle Mead, N.J.)* 2018;47(10):Online. DOI: 10.12788/ajo.2018.0086. [Context Link 1, 2] View abstract...
39. Vajapey SP, Contreras ES, Cvetanovich GL, Neviasser AS. Neurologic complications in primary anatomic and reverse total shoulder arthroplasty: A review. *Journal of Clinical Orthopaedics and Trauma* 2021;20:Online. DOI: 10.1016/j.jcot.2021.06.005. [Context Link 1] View abstract...
40. Burnett Z, Werner BC. Risk factors, management, and prognosis of brachial plexopathy following reverse total shoulder arthroplasty. *Orthopedic Clinics of North America* 2022;53(2):215-221. DOI: 10.1016/j.ocl.2021.11.007. [Context Link 1] View abstract...
41. Florczyński M, Paul R, Leroux T, Baltzer H. Prevention and treatment of nerve injuries in shoulder arthroplasty. *Journal of Bone and Joint Surgery. American Volume* 2021;103(10):935-946. DOI: 10.2106/JBJS.20.01716. [Context Link 1] View abstract...
42. Patel MS, Daher M, Fuller DA, Abboud JA. Incidence, risk factors, prevention, and management of peripheral nerve injuries following shoulder arthroplasty. *Orthopedic Clinics of North America* 2022;53(2):205-213. DOI: 10.1016/j.ocl.2021.11.006. [Context Link 1] View abstract...
43. Kim CY, Sivasundaram L, LaBelle MW, Trivedi NN, Liu RW, Gillespie RJ. Predicting adverse events, length of stay, and discharge disposition following shoulder arthroplasty: a comparison of the Elixhauser Comorbidity Measure and Charlson Comorbidity Index. *Journal of Shoulder and Elbow Surgery* 2018;27(10):1748-1755. DOI: 10.1016/j.jse.2018.03.001. [Context Link 1] View abstract...
44. Wang KY, Quan T, Gu A, Best MJ, Stadecker M, Srikumaran U. Increased severity of anemia is associated with postoperative complications following primary total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2021;30(10):2393-2400. DOI: 10.1016/j.jse.2021.01.022. [Context Link 1] View abstract...
45. Amirian B, et al. A nationwide analysis on the effects of chronic obstructive pulmonary disease following primary total shoulder arthroplasty for glenohumeral osteoarthritis. *Shoulder and Elbow* 2022;14(3):278-285. DOI: 10.1177/1758573221993828. [Context Link 1] View abstract...
46. Burns KA, Robbins LM, LeMarr AR, Fortune K, Morton DJ, Wilson ML. Modifiable risk factors increase length of stay and 90-day cost of care after shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2022;31(1):2-7. DOI: 10.1016/j.jse.2021.08.010. [Context Link 1] View abstract...
47. Xu S, Baker DK, Woods JC, Brabston EW, Ponce BA. Risk factors for earlyReadmission after anatomical or reverse total shoulder arthroplasty. *American Journal of Orthopedics (Belle Mead, N.J.)* 2016;45(6):E386-E392. [Context Link 1, 2] View abstract...
48. Lovy AJ, Keswani A, Beck C, Dowdell JE, Parsons BO. Risk factors for and timing of adverse events after total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2017;26(6):1003-1010. DOI: 10.1016/j.jse.2016.10.019. [Context Link 1, 2] View abstract...
49. Lung BE, Bisogno M, Kanjiya S, Komatsu DE, Wang ED. Early postoperative complications and discharge time in diabetic patients undergoing total shoulder arthroplasty. *Journal of Orthopaedic Surgery and Research* 2019;14(1):9. DOI: 10.1186/s13018-018-1051-3. [Context Link 1, 2] View abstract...
50. Doan MK, et al. Increasing severity of anemia is associated with poorer 30-day outcomes for total shoulder arthroplasty. *JSES International* 2021;5(3):360-364. DOI: 10.1016/j.jseint.2021.02.001. [Context Link 1, 2] View abstract...
51. Aibinder WR, Sanchez-Sotelo J. Venous thromboembolism prophylaxis in shoulder surgery. *Orthopedic Clinics of North America* 2018;49(2):257-263. DOI: 10.1016/j.ocl.2017.11.012. [Context Link 1] View abstract...
52. Cunningham G, et al. A single dose of tranexamic acid reduces blood loss after reverse and anatomic shoulder arthroplasty: a randomized controlled trial. *Journal of Shoulder and Elbow Surgery* 2021;30(7):1553-1560. DOI: 10.1016/j.jse.2020.11.022. [Context Link 1] View abstract...
53. Kennedy JS, et al. The American Society of Shoulder and Elbow Therapists' consensus statement on rehabilitation for anatomic total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2020;29(10):2149-2162. DOI: 10.1016/j.jse.2020.05.019. [Context Link 1, 2] View abstract...
54. MarketScan Database, 2012-2013 (Copyright ©2012-2013 Truven Health Analytics Inc. All Rights Reserved.); proprietary health insurance data sources (2013-2014); and Medicare 100% Standard Analytical File (2013). [Context Link 1]
55. Preyde M, Brassard K. Evidence-based risk factors for adverse health outcomes in older patients after discharge home and assessment tools: a systematic review. *Journal of Evidence-Based Social Work* 2011;8(5):445-68. DOI: 10.1080/15433714.2011.542330. [Context Link 1] View abstract...
56. van Walraven C, et al. Derivation and validation of an index to predict early death or unplanned readmission after discharge from hospital to the community. *Canadian Medical Association Journal* 2010;182(6):551-7. DOI: 10.1503/cmaj.091117. [Context Link 1] View abstract...
57. Hasan O, et al. Hospital readmission in general medicine patients: a prediction model. *Journal of General Internal Medicine* 2010;25(3):211-9. DOI: 10.1007/s11606-009-1196-1. [Context Link 1] View abstract...
58. Billings J, Blunt I, Steventon A, Georghiou T, Lewis G, Bardsley M. Development of a predictive model to identify inpatients at risk of re-admission within 30 days of discharge (PARR-30). *BMJ Open* 2012;2(4):e001667. DOI: 10.1136/bmjopen-2012-001667. [Context Link 1] View abstract...
59. Berkowitz SA, Anderson GF. Medicare beneficiaries most likely to be readmitted. *Journal of Hospital Medicine* 2013;8(11):639-41. DOI: 10.1002/jhm.2074. [Context Link 1] View abstract...
60. Donze J, Lipsitz S, Bates DW, Schnipper JL. Causes and patterns of readmissions in patients with common comorbidities: retrospective cohort study. *British Medical Journal* 2013;347:f7171. [Context Link 1] View abstract...
61. Morris MS, Deierhoi RJ, Richman JS, Altom LK, Hawn MT. The relationship between timing of surgical complications and hospital readmission. *JAMA Surgery* 2014;149(4):348-54. DOI: 10.1001/jamasurg.2013.4064. [Context Link 1] View abstract...
62. Holzgrefe RE, Wilson JM, Staley CA, Anderson TL, Wagner ER, Gottschalk MB. Modified frailty index is an effective risk-stratification tool for patients undergoing total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2019;28(7):1232-1240. DOI: 10.1016/j.jse.2018.12.004. [Context Link 1] View abstract...

63. Cvetanovich GL, et al. Tranexamic acid reduces blood loss after primary shoulder arthroplasty: a double-blind, placebo-controlled, prospective, randomized controlled trial. *Journal of Shoulder and Elbow Surgery Open Access* 2018;2(1):23-27. DOI: 10.1016/j.jses.2018.01.002. [Context Link 1] View abstract...
64. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *New England Journal of Medicine* 2009;360(14):1418-28. DOI: 10.1056/NEJMsa0803563. [Context Link 1, 2, 3] View abstract...
65. Garcia-Perez L, Linertova R, Lorenzo-Riera A, Vazquez-Diaz JR, Duque-Gonzalez B, Sarria-Santamera A. Risk factors for hospital readmissions in elderly patients: a systematic review. *Quarterly Journal of Medicine* 2011;104(8):639-51. DOI: 10.1093/qjmed/hcr070. [Context Link 1, 2] View abstract...
66. Woz S, et al. Gender as risk factor for 30 days post-discharge hospital utilisation: a secondary data analysis. *BMJ Open* 2012;2(2):e000428. DOI: 10.1136/bmjopen-2011-000428. [Context Link 1, 2] View abstract...
67. Jack BW, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. *Annals of Internal Medicine* 2009;150(3):178-87. [Context Link 1] View abstract...
68. Arbaje AI, Wolff JL, Yu Q, Powe NR, Anderson GF, Boulton C. Postdischarge environmental and socioeconomic factors and the likelihood of early hospital readmission among community-dwelling Medicare beneficiaries. *Gerontologist* 2008;48(4):495-504. [Context Link 1] View abstract...
69. Silverstein MD, Qin H, Mercer SQ, Fong J, Haydar Z. Risk factors for 30-day hospital readmission in patients ≥65 years of age. *Proceedings (Baylor University Medical Center)* 2008;21(4):363-72. [Context Link 1] View abstract...
70. Hudson T. The role of social determinates of health in discharge practices. *Nursing Clinics of North America* 2021;56(3):369-378. DOI: 10.1016/j.cnur.2021.04.004. [Context Link 1, 2] View abstract...
71. Mayer RS, Noles A, Vinh D. Determination of postacute hospitalization level of care. *Medical Clinics of North America* 2020;104(2):345-357. DOI: 10.1016/j.mcna.2019.10.011. [Context Link 1] View abstract...
72. Price MC. Musculoskeletal trauma and orthopedic surgery. In: Harding MM, Kwong J, Roberts D, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 11th ed. St. Louis, MO: Mosby; 2020:1444-1476. [Context Link 1]
73. Musculoskeletal disorders. In: Nettina SM, editor. *Lippincott Manual of Nursing Practice*. 11th ed. Philadelphia: Wolters Kluwer Health | Lippincott Williams & Wilkins; 2019:857-892. [Context Link 1]
74. Roles, functions, and preparation of case management team members. In: Powell SK, Tahan H, editors. *Case Management a Practical Guide for Education and Practice*. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:35-60. [Context Link 1]
75. Saleeby J. Communication and collaboration. In: Perry AG, Potter PA, Ostendorf WR, editors. *Nursing Interventions and Clinical Skills*. 7th ed. Elsevier; 2020:9-21. [Context Link 1, 2]
76. National Patient Safety Goals. 2022 National Patient Safety Goals [Internet] Joint Commission on Accreditation of Healthcare Organizations. Accessed at: https://www.jointcommission.org/standards_information/npsgs.aspx. Updated 2022 [accessed 2022 Oct 18] [Context Link 1]
77. The nursing process in drug therapy and patient safety. In: Karch AM, Tucker RG, editors. *Focus on Nursing Pharmacology*. 8th ed. Philadelphia, PA: Wolters Kluwer; 2020:45-55. [Context Link 1]
78. Ostendorf WR. Preparation for safe medication administration. In: Perry AG, Potter PA, Ostendorf WR, editors. *Nursing Interventions and Clinical Skills*. 7th ed. Elsevier; 2020:551-567. [Context Link 1]
79. Transitional planning: understanding levels and transitions of care. In: Powell SK, Tahan H, editors. *Case Management a Practical Guide for Education and Practice*. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:156-211. [Context Link 1]
80. Case management standards and professional organizations. In: Powell SK, Tahan H, editors. *Case Management a Practical Guide for Education and Practice*. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:314-354. [Context Link 1]
81. Charles MD, Cvetanovich G, Sumner-Parilla S, Nicholson GP, Verma N, Romeo AA. Outpatient shoulder arthroplasty: outcomes, complications, and readmissions in 2 outpatient settings. *Journal of Shoulder and Elbow Surgery* 2019;28(6S):S118-S123. DOI: 10.1016/j.jse.2019.04.006. [Context Link 1, 2] View abstract...
82. Ratliff CR. Inflammation and healing. In: Harding MM, Kwong J, Roberts D, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 11th ed. St. Louis, MO: Mosby; 2020:156-174. [Context Link 1]
83. Musculoskeletal care plans. In: Gulanick M, Myers JL, editors. *Nursing Care Plans*. 10th ed. Elsevier; 2022:590-635. [Context Link 1]
84. Wound Care Products. NICE Key Therapeutic Topic KTT14 [Internet] National Institute for Health and Care Excellence. 2019 Sep Accessed at: <https://www.nice.org.uk/guidance>. [accessed 2022 Oct 22] [Context Link 1]
85. Centers for Medicare and Medicaid Services. "Admissions." 42 CFR 412.3 Washington, DC 2023 Jul [accessed 2023 Aug 02] Accessed at: <http://www.gpoaccess.gov/cfr/index.html>. [Context Link 1]
86. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2023 Aug 02] Accessed at: <http://www.gpoaccess.gov/cfr/index.html>. [Context Link 1]
87. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2023 Jul [accessed 2023 Aug 02] Accessed at: <http://www.gpoaccess.gov/cfr/index.html>. [Context Link 1]
88. Centers for Medicare & Medicaid Services. Medicare Benefit Policy Manual. Chapter 1-Inpatient hospital services covered under part A [Internet] Centers for Medicare & Medicaid Services. 2017 Mar10 Accessed at: <http://www.cms.gov/manuals/Downloads/bp102c01.pdf>. [accessed 2017 Oct 04] [Context Link 1]
89. Medicare Benefit Policy Manual. Chapter 6 - hospital services covered under Part B rev. 215 [Internet] Centers for Medicare & Medicaid Services. 2015 Dec Accessed at: <http://www.cms.gov/manuals/>. [accessed 2017 Feb 28] [Context Link 1]

90. Centers for Medicare & Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services [Internet] Centers for Medicare & Medicaid Services. Rev. 11901; 2023 Mar 16 Accessed at: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/>. [accessed 2023 Aug 02] [Context Link 1]
91. Centers for Medicare & Medicaid Services. Medicare Program Integrity Manual. Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment [Internet] Centers for Medicare & Medicaid Services. Rev. 10365; 2020 Oct 02 Accessed at: <https://www.cms.gov/regulations-and-guidance/regulations-and-guidance>. [accessed 2023 Aug 02] [Context Link 1]
92. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx?> Updated 2023 [accessed 2023 Aug 02] [Context Link 1]

Footnotes

[A] See Ambulatory Surgery Discharge and Complications: Common Complications and Conditions [↗](#) ISC for further information. [A in Context Link 1]

[B] See Clinical Indications for Procedure in this guideline. [B in Context Link 1]

[C] Some patients may have their hydration needs met via alternative means (eg, percutaneous endoscopic gastrostomy tube). [C in Context Link 1]

[D] Use Multimodal analgesia or individual analgesic agent as indicated. [D in Context Link 1]

[E] Discharge instructions should be given in the patient's and caregiver's native language using trained language interpreters whenever possible.(70) [E in Context Link 1]

Definitions

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2)(3)(4)(5):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

1. Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:34-41.e1.
2. Lewis J, Patel B. Shock. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 2nd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2018:69-78.
3. Ingbar DH, Thiele H. Cardiogenic shock and pulmonary edema. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2250-2257.
4. Brant EB, Seymour CW, Angus DC. Sepsis and septic shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2241-2249.
5. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.

Hypotension absent

- Hypotension absent, as indicated by **1 or more** of the following(1)(2)(3)(4):
 - SBP greater than or equal to 90 mm Hg in adult or child 10 years or older
 - Mean arterial pressure^[A] greater than or equal to 70 mm Hg in adult or child 10 years or older
 - Mean arterial pressure^[A] at patient's baseline (eg, healthy adult with low SBP), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - SBP greater than or equal to sum of 70 mm Hg plus twice patient's age in years in child 1 to 9 years of age
 - SBP greater than or equal to 70 mm Hg in infant 1 to 11 months of age

References

1. Jones D, Di Francesco L. Hypotension. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:657-64.

2. Massaro AF. Approach to the patient with shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2235-2241.
3. Horeczko T. Pediatric cardiac disorders. In: Walls RM, editor. Rosen's Emergency Medicine. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2109-2131.e1.
4. Singh S, Holmes JF. Pediatric trauma. In: Walls RM, editor. Rosen's Emergency Medicine. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2052-2066.e3.

Footnotes

- A. The mean arterial pressure takes into account both systolic and diastolic blood pressure readings and is calculated as Mean Arterial Pressure (MAP) = 1/3 SBP + 2/3 DBP.

Multimodal analgesia

- Multimodal analgesia involves the utilization of 2 or more analgesic agents with different mechanisms of action in order to provide additive or synergistic pain control, while minimizing side effects and reliance on opioids.(1)(2)(3)

References

1. Hanna MN, Ouanes JP, Tomas VG. Postoperative pain and other acute pain syndromes. In: Benzon HT, Rathmell JP, Wu CL, Turk DC, Argoff CE, Hurley RW, editors. Practical Management of Pain. 5th ed. Philadelphia, PA: Elsevier Mosby; 2014:271-297.e11.
2. George S, Johns M. Review of nonopioid multimodal analgesia for surgical and trauma patients. American Journal of Health-System Pharmacy 2020;77(24):2052-2063. DOI: 10.1093/ajhp/zxaa301.
3. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMra1807061.

Orthostatic hypotension

- Orthostatic hypotension,^{[A][B]} as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:566-571.e1.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Readmission Risk Assessment

- Risk of readmission is increased by presence of **1 or more** of the following(1)(2)(3)(4)(5)(6)(7)(8)(9)(10):
 - Hospitalization (nonelective) in past 6 months(11)(12)(13)
 - 2 or more emergency department visits in past 6 months
 - No source of outpatient care other than emergency department (eg, no primary care provider)(13)(14)
 - Severe care transition barriers (eg, no caregiver, homeless)(11)(12)(15)
 - Severe or end-stage renal disease (on dialysis or GFR less than 30 mL/min/1.73m² (0.5 mL/sec/1.73m²))(11)(16)

 eGFR - Adult Calculator  eGFR - Pediatric Calculator

- Anemia(17)
- Revision shoulder arthroplasty(18)
- Inflammatory arthritis etiology(19)
- Diabetes mellitus necessitating treatment with insulin(20)
- Arthroplasty performed due to traumatic proximal humerus fracture(21)

References

1. MarketScan Database, 2012-2013 (Copyright ©2012-2013 Truven Health Analytics Inc. All Rights Reserved.); proprietary health insurance data sources (2013-2014); and Medicare 100% Standard Analytical File (2013).
2. Preyde M, Brassard K. Evidence-based risk factors for adverse health outcomes in older patients after discharge home and assessment tools: a systematic review. *Journal of Evidence-Based Social Work* 2011;8(5):445-68. DOI: 10.1080/15433714.2011.542330.
3. van Walraven C, et al. Derivation and validation of an index to predict early death or unplanned readmission after discharge from hospital to the community. *Canadian Medical Association Journal* 2010;182(6):551-7. DOI: 10.1503/cmaj.091117.
4. Hasan O, et al. Hospital readmission in general medicine patients: a prediction model. *Journal of General Internal Medicine* 2010;25(3):211-9. DOI: 10.1007/s11606-009-1196-1.
5. Billings J, Blunt I, Steventon A, Georghiou T, Lewis G, Bardsley M. Development of a predictive model to identify inpatients at risk of re-admission within 30 days of discharge (PARR-30). *BMJ Open* 2012;2(4):e001667. DOI: 10.1136/bmjopen-2012-001667.
6. Berkowitz SA, Anderson GF. Medicare beneficiaries most likely to be readmitted. *Journal of Hospital Medicine* 2013;8(11):639-41. DOI: 10.1002/jhm.2074.
7. Donze J, Lipsitz S, Bates DW, Schnipper JL. Causes and patterns of readmissions in patients with common comorbidities: retrospective cohort study. *British Medical Journal* 2013;347:f7171.
8. Morris MS, Deierhoj RJ, Richman JS, Altom LK, Hawn MT. The relationship between timing of surgical complications and hospital readmission. *JAMA Surgery* 2014;149(4):348-54. DOI: 10.1001/jamasurg.2013.4064.
9. Holzgrefe RE, Wilson JM, Staley CA, Anderson TL, Wagner ER, Gottschalk MB. Modified frailty index is an effective risk-stratification tool for patients undergoing total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2019;28(7):1232-1240. DOI: 10.1016/j.jse.2018.12.004.
10. Cvetanovich GL, et al. Tranexamic acid reduces blood loss after primary shoulder arthroplasty: a double-blind, placebo-controlled, prospective, randomized controlled trial. *Journal of Shoulder and Elbow Surgery Open Access* 2018;2(1):23-27. DOI: 10.1016/j.jses.2018.01.002.
11. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *New England Journal of Medicine* 2009;360(14):1418-28. DOI: 10.1056/NEJMsa0803563.
12. Garcia-Perez L, Linertova R, Lorenzo-Riera A, Vazquez-Diaz JR, Duque-Gonzalez B, Sarria-Santamera A. Risk factors for hospital readmissions in elderly patients: a systematic review. *Quarterly Journal of Medicine* 2011;104(8):639-51. DOI: 10.1093/qjmed/hcr070.
13. Woz S, et al. Gender as risk factor for 30 days post-discharge hospital utilisation: a secondary data analysis. *BMJ Open* 2012;2(2):e000428. DOI: 10.1136/bmjopen-2011-000428.
14. Jack BW, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. *Annals of Internal Medicine* 2009;150(3):178-87.
15. Arbaje AI, Wolff JL, Yu Q, Powe NR, Anderson GF, Boult C. Postdischarge environmental and socioeconomic factors and the likelihood of early hospital readmission among community-dwelling Medicare beneficiaries. *Gerontologist* 2008;48(4):495-504.
16. Silverstein MD, Qin H, Mercer SQ, Fong J, Haydar Z. Risk factors for 30-day hospital readmission in patients ≥65 years of age. *Proceedings (Baylor University. Medical Center)* 2008;21(4):363-72.
17. Doan MK, et al. Increasing severity of anemia is associated with poorer 30-day outcomes for total shoulder arthroplasty. *JSES International* 2021;5(3):360-364. DOI: 10.1016/j.jseint.2021.02.001.
18. Xu S, Baker DK, Woods JC, Brabston EW, Ponce BA. Risk factors for early readmission after anatomical or reverse total shoulder arthroplasty. *American Journal of Orthopedics (Belle Mead, N.J.)* 2016;45(6):E386-E392.
19. Lovy AJ, Keswani A, Beck C, Dowdell JE, Parsons BO. Risk factors for and timing of adverse events after total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2017;26(6):1003-1010. DOI: 10.1016/j.jse.2016.10.019.
20. Lung BE, Bisogno M, Kanjiya S, Komatsu DE, Wang ED. Early postoperative complications and discharge time in diabetic patients undergoing total shoulder arthroplasty. *Journal of Orthopaedic Surgery and Research* 2019;14(1):9. DOI: 10.1186/s13018-018-1051-3.
21. Malik AT, Bishop JY, Neviasser AS, Beals CT, Jain N, Khan SN. Shoulder arthroplasty for a fracture is not the same as shoulder arthroplasty for osteoarthritis: implications for a bundled payment model. *Journal of the American Academy of Orthopedic Surgeons* 2019;27(24):927-932. DOI: 10.5435/JAAOS-D-18-00268.

Social Determinants of Health Assessment

- Risk of poor health outcomes may be increased by the presence of **1 or more** of the following social determinants of health(1)(2)(3)(4):
 - Housing insecurity, as indicated by **1 or more** of the following:
 - Individual or caregiver's current living situation is **1 or more** of the following(5):
 - Does not have own housing (eg, staying in a hotel, shelter, or with others)

- Has own housing (eg, house, apartment), but at risk of losing it in the future (ie, behind on rent or mortgage)
- Has own housing (eg, house, apartment), but has lived in 3 or more places in past year
- Current housing has **1 or more** of the following:
 - Electrical appliances (eg, stove, refrigerator) not working or unavailable
 - Insufficient heating or cooling
 - Insufficient ventilation
 - Lead paint or pipes
 - Mold
 - Pests (eg, bugs) or rodents
 - Smoke detectors not working or unavailable
- Food insecurity, as indicated by **1 or more** of the following(6):
 - In the past year, individual or caregiver ran out of food and did not have money to buy more food.
 - In the past year, individual or caregiver worried that they would run out of food before they received money to buy more food.
- Insufficient transportation, as indicated by **1 or more** of the following(7):
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of transportation.
 - In the past year, individual or caregiver missed nonmedical activities, work, or could not get things needed for daily living due to lack of transportation.
- Insufficient utilities, as indicated by **1 or more** of the following(8):
 - Utilities (eg, electricity, water, gas, or oil) are currently shut off or unavailable.
 - In the past year, electric, water, gas, or oil company threatened to shut off services.
- Personal safety risk, as indicated by **2 or more** of the following(6):
 - Individual is sometimes or frequently physically hurt by another person (including family member).
 - Individual is sometimes or frequently insulted or talked down to by another person (including family member).
 - Individual is sometimes or frequently threatened with physical harm by another person (including family member).
 - Individual is sometimes or frequently screamed or cursed at by another person (including family member).
- Insufficient dependent care, as indicated by **1 or more** of the following:
 - In the past year, individual or caregiver was unable to work due to lack of dependent care.
 - In the past year, individual or caregiver was unable to work more (additional) hours due to lack of dependent care.
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of dependent care.
 - In the past year, individual or caregiver missed nonmedical activities (eg, school, church, social activity) due to lack of dependent care.
- Depression risk, as indicated by **ALL** of the following:
 - In the past 2 weeks, individual had little interest or pleasure in normal activities on at least several days.
 - In the past 2 weeks, individual felt down, depressed, or hopeless on at least several days.

References

1. Social Determinants of Health. [Internet] World Health Organization. Accessed at: https://www.who.int/social_determinants/sdh_definition/en/. Updated 2022 [accessed 2022 Apr 20]
2. Moen M, Storr C, German D, Friedmann E, Johantgen M. A review of tools to screen for social determinants of health in the United States: a practice brief. *Population Health Management* 2020;23(6):422-429. DOI: 10.1089/pop.2019.0158.
3. Daniel-Robinson L, Moore JE. Innovation and Opportunities to Address Social Determinants of Health in Medicaid Managed Care. [Internet] Institute for Medicaid Innovation. 2019 Jan Accessed at: <https://www.medicaidinnovation.org/>. [accessed 2022 Oct 18]
4. Billioux A, Verlander K, Anthony S, Alley D. Standardized Screening for Health-Related Social Needs in Clinical Settings: the Accountable Health Communities Screening Tool. [Internet] National Academy of Sciences. 2017 May Accessed at: <https://nam.edu/>. [accessed 2022 Sep 14]
5. Sandel M, et al. Unstable housing and caregiver and child health in renter families. *Pediatrics* 2018;14(2):e20172199. DOI: 10.1542/peds.2017-2199.
6. Children's HealthWatch Survey. Screening Instrument [Internet] Children's HealthWatch. 2020 Sep Accessed at: <https://childrenshealthwatch.org/>. [accessed 2022 Oct 27]
7. PRAPARE®: Protocol for Responding to and Assessing Patient Assets, Risks, and Experiences Screening Tool. [Internet] Association of Asian Pacific Community Health Organizations (AAPCHO) and National Association of Community Health Centers (NACHC). 2016 Sep Accessed at: <https://prapare.org/the-prapare-screening-tool/>. [accessed 2022 Sep 26]
8. Cook JT, et al. A brief indicator of household energy security: associations with food security, child health, and child development in US infants and toddlers. *Pediatrics* 2008;122(4):e867-75. DOI: 10.1542/peds.2008-0286.

Tachycardia absent

- Tachycardia absent, as indicated by **1 or more** of the following(1)(2):
 - Heart rate less than or equal to 100 beats per minute in adult or child 6 years or older
 - Heart rate less than or equal to 115 beats per minute in child 3 to 5 years of age

- Heart rate less than or equal to 125 beats per minute in child 1 or 2 years of age
- Heart rate less than or equal to 130 beats per minute in infant 6 to 11 months of age
- Heart rate less than or equal to 150 beats per minute in infant 3 to 5 months of age
- Heart rate less than or equal to 160 beats per minute in infant 1 or 2 months of age

References

1. Southmayd GL. Tachycardia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:729-739.
2. Pediatric parameters and equipment. In: Kleinman K, McDaniel L, Molloy M, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 22nd ed. 202: Elsevier; 2021:frontpiece tables.

Codes

ICD-10 Diagnosis: M05.011, M05.012, M05.019, M05.611, M05.612, M05.619, M05.711, M05.712, M05.719, M05.811, M05.812, M05.819, M06.011, M06.012, M06.019, M12.511, M12.512, M12.519, M12.811, M12.812, M12.819, M12.9, M13.111, M13.112, M13.119, M19.011, M19.012, M19.019, M19.112, M19.119, M19.212, M19.219, M80.011A, M80.012A, M80.019A, M80.811A, M80.812A, M80.819A, M84.311A, M84.312A, M84.319A, M84.411A, M84.412A, M84.419A, M84.511A, M84.512A, M84.519A, M84.611A, M84.612A, M84.619A, M87.011, M87.012, M87.019, M87.021, M87.022, M87.029, M87.121, M87.122, M87.129, M87.221, M87.222, M87.229, M87.321, M87.322, M87.329, M87.821, M87.822, M87.829, M90.511, M90.512, M90.519, M96.621, M96.622, M96.629, M97.31XA, M97.32XA, S42.001A, S42.001B, S42.002A, S42.002B, S42.009A, S42.009B, S42.031A, S42.031B, S42.032A, S42.032B, S42.033A, S42.033B, S42.034A, S42.034B, S42.035A, S42.035B, S42.036A, S42.036B, S42.121A, S42.121B, S42.122A, S42.122B, S42.123A, S42.123B, S42.124A, S42.124B, S42.125A, S42.125B, S42.126A, S42.126B, S42.131A, S42.131B, S42.132A, S42.132B, S42.133A, S42.133B, S42.134A, S42.134B, S42.135A, S42.135B, S42.136A, S42.136B, S42.141A, S42.141B, S42.142A, S42.142B, S42.143A, S42.143B, S42.144A, S42.144B, S42.145A, S42.145B, S42.146A, S42.146B, S42.151A, S42.151B, S42.152A, S42.152B, S42.153A, S42.153B, S42.154A, S42.154B, S42.155A, S42.155B, S42.156A, S42.156B, S42.201A, S42.201B, S42.202A, S42.202B, S42.209A, S42.209B, S42.211A, S42.211B, S42.212A, S42.212B, S42.213A, S42.213B, S42.214A, S42.214B, S42.215A, S42.215B, S42.216A, S42.216B, S42.221A, S42.221B, S42.222A, S42.222B, S42.223A, S42.223B, S42.224A, S42.224B, S42.225A, S42.225B, S42.226A, S42.226B, S42.231A, S42.231B, S42.232A, S42.232B, S42.239A, S42.239B, S42.241A, S42.241B, S42.242A, S42.242B, S42.249A, S42.249B, S42.251A, S42.251B, S42.252A, S42.252B, S42.253A, S42.253B, S42.254A, S42.254B, S42.255A, S42.255B, S42.256A, S42.256B, S42.261A, S42.261B, S42.262A, S42.262B, S42.263A, S42.263B, S42.264A, S42.264B, S42.265A, S42.265B, S42.266A, S42.266B, S42.271A, S42.272A, S42.279A, S42.291A, S42.291B, S42.292A, S42.292B, S42.293A, S42.293B, S42.294A, S42.294B, S42.295A, S42.295B, S42.296A, S42.296B, S42.90XA, S42.90XB, S42.91XA, S42.91XB, S42.92XA, S42.92XB, S49.001A, S49.002A, S49.009A, S49.011A, S49.012A, S49.019A, S49.021A, S49.022A, S49.029A, S49.031A, S49.032A, S49.039A, S49.041A, S49.042A, S49.049A, S49.091A, S49.092A, S49.099A, T84.018A, T84.028A, T84.038A, T84.058A, T84.098A, T84.110A, T84.111A, T84.120A, T84.121A, T84.190A, T84.191A, T84.318A, T84.328A, T84.398A, T84.410A, T84.418A, T84.420A, T84.428A, T84.490A, T84.498A, T84.59XA, T84.84XA, T84.89XA, T84.9XXA [Hide]

ICD-10 Procedure: 0RRJ00Z, 0RRJ07Z, 0RRJ0JZ, 0RRJ0KZ, 0RRK00Z, 0RRK07Z, 0RRK0JZ, 0RRK0KZ

CPT®: 23472, 23473, 23474

CPT copyright 2022 American Medical Association. All rights reserved.

MCG Health
 Inpatient & Surgical Care 27th Edition
 Copyright © 2023 MCG Health, LLC
 All Rights Reserved

Last Update: 9/21/2023 11:27:09 AM
 Build Number: 27.2.2023092114759.013030